This listing of claims will replace all prior versions and listings of claims in the application.

Listing of Claims

1. (Previously Presented) An acylated indanyl amine according to the general formula (I) in any of its stereoisomeric forms or a mixture thereof in any ratio or a pharmaceutically acceptable salt thereof

wherein

R¹ and R⁴ are independently from each other selected from the group consisting of:

H;

unsubstituted and at least monosubstituted C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl and C₂-C₁₀-alkynyl, the substituents of which are selected from the group consisting of F, OH, C₁-C₈-alkoxy, (C₁-C₈-alkyl)mercapto, CN, COOR⁶, CONR⁷R⁸, and unsubstituted and at least monosubstituted phenyl, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy and CF₃;

unsubstituted and at least monosubstituted phenyl, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy and CF₃;

R⁹CO; CONR¹⁰R¹¹; COOR¹²; CF₃; halogens; pseudohalogens; NR¹³R¹⁴; OR¹⁵; S(O)_mR¹⁶; SO₂NR¹⁷R¹⁸; and NO₂;

R² and R³ are independently from each other selected from the group consisting of:

H;

halogens;

pseudohalogens;

unsubstituted and at least monosubstituted C_1 - C_{10} -alkyl the substituents of which are selected from the group consisting of OH and phenyl;

OH;

 C_1 - C_{10} -alkoxy;

phenoxy;

 $S(O)_{m}R^{19};$

CF₃:

CN;

NO₂;

(C₁-C₁₀-alkyl)amino;

di(C₁-C₁₀-alkyl)amino;

(C₁-C₆-alkyl)-CONH-;

unsubstituted and at least monosubstituted phenyl-CONH- and phenyl-SO₂-O-, the substituents of which are selected from the group consisting of halogens, pseudohalogens, CH₃ and methoxy;

```
(C_1-C_6-alkyl)SO_2-O_-;
```

unsubstituted and at least monosubstituted (C_1 - C_6 -alkyl)CO, the substituents of which are selected from the group consisting of F and di(C_1 - C_3 -alkyl)amino;

and phenyl-CO, the phenyl part of which can be substituted by one or more substituents from the group consisting of C₁-C₃-alkyl, halogens and methoxy;

A is selected from the group consisting of CH2, CHOH and CH-(C1-C3-alkyl);

B is selected from the group consisting of CH₂ and CH-(C₁-C₃-alkyl);

R⁵ is a benzo[1,3] dioxole group optionally substituted with one or more substituents selected from the group consisting of:

halogens;

pseudohalogens;

NH₂;

unsubstituted and at least monosubstituted C_1 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_2 - C_{10} -alkynyl, C_1 - C_{10} -alkoxy, $(C_1$ - C_{10} -alkyl)amino, and $di(C_1$ - C_{10} -alkyl)amino, the substituents of which are selected from the group consisting of F, OH, C_1 - C_8 -alkoxy, aryloxy, $(C_1$ - C_8 -alkyl)mercapto, NH₂, $(C_1$ - C_8 -alkyl)amino, and $di(C_1$ - C_8 -alkyl)amino; C_3 - C_5 -alkandiyl;

```
phenyl;
aryl-substituted C1-C4-alkyl;
CF<sub>3</sub>;
NO<sub>2</sub>;
OH:
phenoxy;
benzyloxy;
(C_1-C_{10}-alkyl)COO;
S(O)_m R^{20};
SH;
phenylamino;
benzylamino;
(C_1-C_{10}-alkyl)-CONH-,
(C_1-C_{10}-alkyl)-CON(C_1-C_4-alkyl)-;
phenyl-CONH-;
phenyl-CON(C_1-C_4-alkyl)-;
(C_1-C_{10}-alkyl)-CO;
phenyl-CO;
CF<sub>3</sub>-CO;
-OCH<sub>2</sub>O-;
-OCF<sub>2</sub>O-;
-OCH2CH2O-;
-CH<sub>2</sub>CH<sub>2</sub>O-;
COOR<sup>21</sup>:
CONR<sup>22</sup>R<sup>23</sup>;
CNH(NH<sub>2</sub>);
SO2NR24R25;
R<sup>26</sup>SO<sub>2</sub>NH-;
R^{27}SO_2N(C_1-C_6-alkyl)-;
```

and wherein all aryl, phenyl, aryl-containing, and phenyl-containing groups, which are optionally present in the said substituents of the benzo[1,3]dioxole group can be substituted by one or more substituents selected from the group consisting of halogens, pseudohalogens, C_1 - C_3 -alkyl, OH, C_1 - C_3 -alkoxy, and CF_3 ;

R⁶ is selected from the group consisting of:

H;

 C_1 - C_{10} -alkyl, which can be substituted by one or more substituents selected from the group consisting of F, C_1 - C_8 -alkoxy, and di(C_1 - C_8 -alkyl)amino;

4 of 11

aryl- $(C_1-C_4$ -alkyl) optionally substituted by one or more substituents selected from the group consisting of halogens, C_1-C_4 -alkoxy, and di $(C_1-C_6$ -alkyl)amino;

R⁷ is selected from the group consisting of:

H.

 C_1 - C_{10} -alkyl which can be substituted by one or more substituents selected from the group consisting of F, C_1 - C_8 -alkoxy, di(C_1 - C_8 -alkyl)amino and phenyl;

phenyl; and indanyl;

and wherein each of the aforementioned aromatic groups can be unsubstituted or carry one or more substituents from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy and CF₃;

 \mathbb{R}^8 is H or \mathbb{C}_1 - \mathbb{C}_{10} -alkyl;

R⁹ is selected from the group consisting of:

 C_1 - C_{10} -alkyl which can be unsubstituted or carry one or more substituents from the group consisting of: F, $(C_1$ - $C_4)$ -alkoxy, di $(C_1$ - C_3 -alkyl)amino;

and unsubstituted and at least monosubstituted phenyl, the substituents of which are selected from the group consisting of C₁-C₃-alkyl, C₁-C₃-alkoxy, halogens, pseudohalogens, and CF₃;

 R^{10} independently has the same meaning as R^7 ;

 R^{11} independently has the same meaning as R^8 ;

R¹² independently has the same meaning as R⁶;

R¹³ is selected from the group consisting of:

H;

C₁-C₆-alkyl;

unsubstituted and substituted phenyl, benzyl, $(C_1-C_6-alkyl)-CO$, and phenyl-CO, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C_1 - C_3 -alkyl, C_1-C_3 -alkoxy, and CF_3 ,

and wherein one or more of these substituents can be present;

R¹⁴ independently has the same meaning as R¹³;

R¹⁵ is selected from the group consisting of:

H:

C₁-C₁₀-alkyl;

 $(C_1-C_3-alkoxy)-C_1-C_3-alkyl;$

and substituted and unsubstituted benzyl, and phenyl, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy, and CF₃, and wherein one or more of these substituents can be present;

R¹⁶ is selected from the group consisting of:

 C_1 - C_{10} -alkyl which can be substituted by one or more substituents selected from the group consisting of F, OH, C_1 - C_8 -alkoxy, aryloxy, $(C_1$ - C_8 -alkyl)mercapto, $(C_1$ - C_8 -alkyl)amino and di $(C_1$ - C_8 -alkyl)amino;

CF₃;

and substituted and unsubstituted phenyl, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy and CF₃, and wherein one or more of these substitutents can be present;

R¹⁷ independently has the same meaning as R⁷;

R¹⁸ independently has the same meaning as R⁸;

R¹⁹ independently has the same meaning as R¹⁶;

R²⁰ independently has the same meaning as R¹⁶;

R²¹ independently has the same meaning as R⁶;

 \mathbb{R}^{22} independently has the same meaning as \mathbb{R}^7 ;

R²³ independently has the same meaning as R⁸;

 R^{24} independently has the same meaning as R^7 ;

R²⁵ independently has the same meaning as R⁸;

R²⁶ independently has the same meaning as R¹⁶;

R²⁷ independently has the same meaning as R¹⁶;

aryl is phenyl, naphth-1-yl or naphth-2-yl; and m is 0, 1 or 2.

2. (Currently Amended) An acylated indanyl amine in any of its stereoisomeric forms or a mixture thereof in any ratio or a pharmaceutically acceptable salt thereof according to claim 1, wherein in the formula (I):

R¹ is selected from the group consisting of: H; C₁-C₄-alkyl; C₁-C₄-alkoxy; CF₂; halogens; pseudohalogens; (C₁-C₄-alkyl)-S(O)_m-; and unsubstituted and at least monosubstituted phenyl, the substituents of which are selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, C₁-C₃-alkoxy, and CF₃;

 R^2 and R^3 are independently from each other selected from the group consisting of: H; halogens; pseudohalogens; and C_1 - C_3 -alkyl;

R⁴ independently has the same meaning as R¹;

A is selected from the group consisting of CH2 and CHOH;

B is selected from the group consisting of CH2 and CH-CH3;

R⁵ is a benzo[1,3] dioxole group optionally substituted with one or more substituents selected from the group consisting of: halogens; CN; NH2; unsubstituted and at least monosubstituted C1-C8-alkyl, C2-C8-alkenyl, C2-C8-alkynyl, C1-C8-alkoxy, (C1-C8alkyl)amino, and di(C1-C8-alkyl)amino, the substituents of which are selected from the group consisting of F, C1-C6-alkoxy, phenoxy, (C1-C6-alkyl)mercapto, NH2, (C1-C6-alkyl)amino, and di(C1-C6-alkyl)amino; C3-C5-alkandiyl; phenyl; phenyl-substituted C1-C2-alkyl; CF3; OH; phenoxy; benzyloxy; $(C_1-C_6-alkyl)COO$; $S(O)_m(C_1-C_6)-alkyl$; $S(O)_m-phenyl$; SH; phenylamino; benzylamino; (C1-C6-alkyl)-CONH-; (C1-C6-alkyl)-CON(C1-C4-alkyl)-; phenyl-CONH-; phenyl-CON(C₁-C₄-alkyl)-; (C₁-C₆-alkyl)-CO; phenyl-CO; CF₃-CO; -OCH₂O-; - $OCF_2O-; -OCH_2CH_2O-, -CH_2CH_2O-; COO(C_1-C_6-alkyl); -CONH_2; -CONH(C_1-C_6-alkyl); -CONH_2(C_1-C_6-alkyl); -CONH_2(C_1-C$ $CON(di(C_1-C_6-alkyl)); CNH(NH_2); -SO_2NH_2; -SO_2NH(C_1-C_6-alkyl); -SO_2NH(phenyl); -S$ $SO_2N(di(C_1-C_6-alkyl)); (C_1-C_6-alkyl)SO_2NH-; (C_1-C_6-alkyl)SO_2N(C_1-C_6-alkyl)-; phenyl-phe$ SO₂NH-; and phenyl-SO₂N(C₁-C₆-alkyl)-; and wherein all phenyl and phenyl-containing groups, which are optionally present in the said substituents of the benzo[1,3] dioxole group, can be substituted by one or more substituents selected from the group consisting of halogens, pseudohalogens, C1-C3-alkyl, OH, C1-C3-alkoxy, and CF3; and

m is 0 or 2.

3. (Currently Amended) An acylated indanyl amine in any of its stereoisomeric forms or a mixture thereof in any ratio or a pharmaceutically acceptable salt thereof according to claim 1, wherein in the formula (I):

R1 is H, halogen or C1-C4-alkyl;

R² and R³ are each H;

R4 independently has the same meaning as R1;

A is CH2;

B is CH₂;

R⁵ is a benzo[1,3] dioxole group optionally substituted with one or more substituents selected from the group consisting of: halogens; CN; NH₂; unsubstituted and at least monosubstituted C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₃-alkoxy, (C₁-C₄-alkyl)amino, and di(C₁-C₄-alkyl)amino, the substituents of which are selected from the group consisting of F, C₁-C₃-alkoxy, (C₁-C₃-alkyl)mercapto, and NH₂; C₃-C₅-alkandiyl; phenyl; phenyl-substituted C₁-C₂-alkyl; CF₃; OH; (C₁-C₄-alkyl)COO; S(O)_m (C₁-C₄)-alkyl; (C₁-C₄-alkyl)-CONH-; (C₁-C₄-alkyl)-CON(C₁-C₄-alkyl)-; (C₁-C₄-alkyl)-CO; phenyl-CO; CF₃-CO; -OCH₂O-; -OCF₂O-; -OCH₂CH₂O-; -CH₂CH₂O-; COO(C₁-C₆-alkyl); -CONH₂; -CONH(C₁-C₄-alkyl); -SO₂NH₂; -SO₂NH₂; -SO₂NH₂; -CONH(C₁-C₄-alkyl); -SO₂NH(phenyl); -SO₂N(di(C₁-C₄-alkyl)); (C₁-C₄-alkyl)SO₂NH-; and (C₁-C₄-alkyl)SO₂N(C₁-C₄-alkyl)-; and wherein all phenyl and phenyl-containing groups, which are optionally present in the said substituents of the benzo[1,3] dioxole group; can be substituted by one or more substituents selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, OH, C₁-C₃-alkoxy, and CF₃; and

m is 0 or 2.

4. (Currently Amended) An acylated indanyl amine in any of its stereoisomeric forms or a mixture thereof in any ratio or a pharmaceutically acceptable salt thereof according to claim 1, wherein in the formula (I)

R¹ is H, halogen or C₁-C₄-alkyl;

R² and R³ are each H;

R4 independently has the same meaning as R1;

A and B are each CH2;

R⁵ is benzo[1,3] dioxole group optionally substituted with one or more substituents selected from the group consisting of: F; Cl; Br; C₁-C₃-alkyl; C₁-C₃-alkoxymethyl; 2-amino-

3,3,3-trifluoro-propyl-; CF₃; C₃-C₅-alkandiyl; phenyl; benzyl; OH; C₁-C₃-alkoxy; phenoxy; trifluoromethoxy; 2,2,2-trifluoroethoxy; (C₁-C₄-alkyl)COO; (C₁-C₃-alkyl)mercapto; phenylmercapto; (C₁-C₃-alkyl)sulfonyl; phenylsulfonyl; NH₂; (C₁-C₄-alkyl)amino; di(C₁-C₄-alkyl)amino; (C₁-C₃-alkyl)-CONH-; (C₁-C₃-alkyl)-SO₂NH-; (C₁-C₃-alkyl)-CO; phenyl-CO; -OCH₂O-; -OCF₂O-; -CH₂CH₂O-; COO(C₁-C₄-alkyl); -CONH₂; -CONH(C₁-C₄-alkyl); -CON(di(C₁-C₄-alkyl)); CN; -SO₂NH₂; -SO₂NH(C₁-C₄-alkyl); and -SO₂N(di(C₁-C₄-alkyl)); and wherein all phenyl and phenyl-containing groups which are optionally present in said benzo[1,3] dioxole group can be substituted by one or more substituents selected from the group consisting of halogens, pseudohalogens, C₁-C₃-alkyl, OH, C₁-C₃-alkoxy, and CF₃.

5. (Previously Presented) An acylated indanyl amine in any of its stereoisomeric forms or a mixture thereof in any ratio or a pharmaceutically acceptable salt thereof according to claim 1, wherein in the formula (I):

R¹ is H, halogen or C₁-C₄-alkyl;

 R^2 , R^3 and R^4 are each H;

A and B are each CH2;

R⁵ is selected from the group consisting of: benzo[1,3]dioxol-5-yl, and 2,2-difluoro-benzo[1,3]dioxol-5-yl.

6. (Previously Presented) An acylated indanyl amine or a pharmaceutically acceptable salt thereof according to claim 1, which is 2,2-diffuoro-benzo[1,3]dioxole-5-carboxylic acid indan-2-ylamide.

7-20. (Canceled)

- 21. (Original) A pharmaceutical preparation comprising an effective dose of at least one compound of the formula (I) as defined in claim 1 in any of its stereoisomeric forms or a mixture thereof in any ratio and/or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.
- 22. (Original) A pharmaceutical preparation according to claim 21, which pharmaceutical preparation is in the form of a pill, tablet, lacquered tablet, sugar-coated tablet, granule, hard or soft gelatin capsule, aqueous, alcoholic or oily solution, syrup, emulsion or

suspension, suppository, solution for injection or infusion, ointment, tincture, spray, transdermal therapeutic systems, nasal spray, aerosol mixture, microcapsule, implant or rod.

23. (Canceled)

- 24. (Previously Presented) The acylated indanyl amine according to claim 1 selected from the group consisting of benzo[1,3]dioxol-5-carboxylic-acid (5-nitro-indan-2-yl)-amide, benzo[1,3]dioxol-5-carboxylic-acid (6-chlor-1-hydroxy-indan-2-yl)-amide, 2,2-difluoro-benzo[1,3]dioxol-5-carboxylic acid indan-2-ylamide, and benzo[1,3]dioxol-5-carboxylic acid indan 2-yl-amide.
- 25. (Previously Presented) The acylated indanyl amine according to claim 24, which is 2,2-difluoro-benzo[1,3]dioxol-5-carboxylic acid indan-2-ylamide.
- 26. (Previously Presented) A pharmaceutical preparation comprising an effective dose of at least one compound of claim 24 and a pharmaceutically acceptable carrier.
- 27. (Previously Presented) A pharmaceutical preparation according to claim 26, which pharmaceutical preparation is in the form of a pill, tablet, lacquered tablet, sugar-coated tablet, granule, hard or soft gelatin capsule, aqueous, alcoholic or oily solution, syrup, emulsion or suspension, suppository, solution for injection or infusion, ointment, tincture, spray, transdermal therapeutic systems, nasal spray, aerosol mixture, microcapsule, implant or rod.